



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of: Stoynova et al.	
	Art Unit: [to be assigned]
Application No.: 10/654898	Examiner: [to be assigned]
Filing Date: September 5, 2003	Atty. Docket: US-100
Title: Method for Producing L-Amino Acid Using Bacterium Belonging to the Genus <i>Escherichia</i> , Lacking Active <i>mlc</i> Gene	

CLAIM FOR PRIORITY UNDER 35 U.S.C. § 119 IN UTILITY APPLICATION

Commissioner of Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

Priority under 35 U.S.C. § 119 is hereby claimed to the following priority document(s), filed in a foreign country within one (1) year prior to the filing of the above-referenced United States utility patent application (35 U.S.C. § 172):

Country	Priority Document Appl. No.	Filing Date
Russia	2002123822	September 6, 2002

A certified copy of the listed priority document and a translation of the front page is submitted herewith. Prompt acknowledgment of this claim and submission is respectfully requested.

Respectfully submitted,

Shelly Guest Cermak
Reg. No. 39,571

Date: January 6, 2004
PTO Customer Number: **000038108**
Ajinomoto U.S.A., Inc.
1120 Connecticut Avenue, Ste. 1010
Washington, D.C. 20036
202.457.0284
202.457.0107 (fax)



Russian agency
for patents and trade marks
(ROSPATENT)

**Federal Institute
of Industrial Property**

Berezhkovskaya nab., 30, corp. 1, Moscow, G-59, GSP-5, 123995

Phone 240 60 15. Telex 114818 PDCh. Fax 243 33 37

Reg.№ 20/12-400

July 23, 2003

CERTIFICATE

Federal Institute of Industrial Property (hereinafter referred as - Institute) do hereby certify, that attached materials are exact copy of the original specification, claims, abstract and drawings (if exist) of the application №2002123822 for issuing the patent for invention, filed to the Institute on September 6th, 2002 (06.09.2002)

Title of invention: Method for producing L-amino acids using bacterium, belonging to the genus *Escherichia*, lacking active *mlc* gene.

Applicant: Closed Joint Stock Company
"Ajinomoto-Genetika Research Institute"

Actual author(s): STOYNOVA Natalia Viktorovna
SYCHEVA Elena Viktorovna
SKOROKHODOVA Alexandra Yurievna
KOZLOV Yuri Ivanovich

SEAL

Head of Department 20 (Signature) A.L.Zhuravlev

РОССИЙСКОЕ АГЕНТСТВО
ПО ПАТЕНТАМ И ТОВАРНЫМ ЗНАКАМ
(РОСПАТЕНТ)



**ФЕДЕРАЛЬНЫЙ ИНСТИТУТ
ПРОМЫШЛЕННОЙ СОБСТВЕННОСТИ**

Бережковская наб., 30, корп. 1, Москва, Г-59, ГСП-5, 123995
Телефон 240 60 15. Телекс 114818 ПДЧ. Факс 243 33 37

Наш № 20/12-400

«23» июля 2003 г.

СПРАВКА

Федеральный институт промышленной собственности (далее – Институт) настоящим удостоверяет, что приложенные материалы являются точным воспроизведением первоначального описания, формулы, реферата и чертежей (если имеются) заявки №2002123822 на выдачу патента на изобретение, поданной в Институт в сентябре месяце 06 дня 2002 года (06.09.2002).

Название изобретения:

Способ получения L- аминокислот с использованием бактерий, принадлежащих к роду Escherichia, содержащих неактивный ген mlc.

Заявитель:

Закрытое акционерное общество
«Научно-исследовательский институт
Аджиномото - Генетика»

Действительные авторы:

СТОЙНОВА Наталия Викторовна
СЫЧЕВА Елена Викторовна
СКОРОХОДОВА Александра Юрьевна
КОЗЛОВ Юрий Иванович



Заведующий отделом 20

А.Л.Журавлев

METHOD FOR PRODUCING L-AMINO ACID USING BACTERIUM, BELONGING TO THE GENUS *ESCHERICHIA*, LACKING ACTIVE *MLC* GENE.

Background of the Invention

Field of invention

The present invention relates to the microbiological industry, specifically to a method for producing L-amino acid using bacterium, belonging to the genus *Escherichia*, wherein *mlc* gene is inactivated.

Description of the related art

Mlc protein is a global regulator (repressor) of carbohydrate metabolism (Decker et al, Mol Microbiol 1998, 27:2:381-90; Kimata et al, Mol Microbiol, 1998, 29:6:1509-19; Plumbridge, Mol Microbiol, 1998, 27:2:369-80). It was shown Mlc protein regulates expression of several genes and operons. There are *ptsG* gene (Kimata et al, Mol Microbiol, 1998, 29:6:1509-19; Plumbridge, Mol Microbiol, 1998, 29:4:1053-63; Kim et al, J Biol Chem, 1999, 274:36:25398-402; Plumbridge, Mol Microbiol 1999, 33:2:260-73; Tanaka et al, Genes Cells, 1999, 4:7:391-9) encoding membrane-bound subunit, IICB(Glc), of glucose phosphotransferase system (PTS), *ptsHIcrr* operon (encoding general PTS proteins) Kimata et al, Mol Microbiol, 1998, 29:6:1509-19; Plumbridge, Mol Microbiol, 1998, 29:4:1053-63; Kim et al, J Biol Chem, 1999, 274:36:25398-402; Plumbridge, Mol Microbiol 1999, 33:2:260-73; Tanaka et al, Genes Cells, 1999, 4:7:391-9), *manXYZ* operon encoding enzyme II of mannose PTS (Plumbridge, Mol Microbiol, 1998, 29:4:1053-63), *malT* gene encoding the activator of maltose regulon (Decker et al, Mol Microbiol 1998, 27:2:381-90).

Genes of *mlc* regulon are also positively regulated by CRP-cAMP complex (Chapon and Colb, J. Bacteriol., 1983, 156:1135-43; Decker et al, Mol. Microbiol., 1998, 27:2:381-90; Kimata et al, Mol. Microbiol., 1998, 29:6:1509-19; Plumbridge, Mol. Microbiol., 1998, 27:2:369-80; Plumbridge, Mol. Microbiol., 1998, 29:4:1053-63; Kim et al, J. Biol. Chem, 1999,

274:36:25398-402; Plumbridge, Mol. Microbiol 1999, 33:2:260-73; Tanaka et al, Genes Cells, 1999, 4:7:391-9)

Regulation of *mlc* gene transcription is remarkably complicated. First, it is negatively regulated by Mlc protein itself. Unphosphorylated EIICB(Glc) (product of *ptsG* gene) can sequester Mlc protein from its binding site by direct protein-protein interaction and therefore induce expression of *mlc* regulon in response of glucose (Tanaka et al, EMBO J, 2000, 19:20, 5344-52; Lee et al, EMBO J, 2000, 19:20:5353-61; Nam et al, EMBO J, 2001, 20:3:491-8). Second, transcription of *mlc* gene is performed by two promoters P1 and P2 (Shin et al, J. Biol. Chem. 2001, 276:28:25871-75). Promoter P1 is recognized only by RNA polymerase, containing the housekeeping sigma factor σ^{70} ($E\sigma^{70}$), while the promoter P2 can be recognized by both $E\sigma^{70}$ and $E\sigma^{32}$, containing the heat shock sigma factor. Thus *mlc* gene belongs to a class of genes, transcribed from the multiple promoters including one recognized by RNA polymerase associated with the alternative sigma factor in order to respond to various environmental conditions. In addition, a highly conserved CRP-binding site present within the *mlc* promoter (Shin et al, J. Biol. Chem. 2001, 276:28:25871-75).

Transport of carbohydrates provided by PTS may be the limiting step in overproduction of some amino acids. So, the inactivation of product of *mlc* gene, which negatively regulates PTS genes expression, seems to be necessary for increasing of amino acid production. In the populations of *Escherichia coli* growing in the glucose-limited environment the polygenic mutations in *mgl*, *mlc* and *malT* genes were found (Manch K., Genetics, 1999, 153:1:5-12). But at the present time there is no report of inactivation of *mlc* gene used for producing of L-amino acid.

Disclosure of the Invention

An object of present invention is to enhance the productivity of L-amino acid producing strains and to provide a method for producing L-amino acid using these strains.

This aim was achieved by finding that the inactivation of *mlc* gene encoding the repressor of carbohydrate metabolism can enhance production of L-amino acid, such as L-threonine. Thus the present invention has been completed.

The present inventions are as follows:

- 1). An L-amino acid producing bacterium belonging to the genus *Escherichia*, wherein the bacterium has been modified to have *mlc* gene inactivated.
- 2). The L-amino acid producing bacterium, wherein L-amino acid is L-threonine.
- 3). A method for producing L-amino acid, which method comprises the steps of:
 - cultivating the bacterium according to any 1 or 2 in a medium to produce and accumulate L-amino acid in the medium, and
 - collecting L-amino acid from the medium.
- 4). The method according to 3), wherein L-amino acid is L-threonine.
- 5). The method according to 4), wherein the bacterium has been modified to have enhanced expression of L-threonine operon.

The present invention is described in details below.

1. Bacterium of the present invention

The bacterium of the present invention is an L-amino acid producing bacterium belonging to the genus *Escherichia*, wherein the bacterium has been modified to have *mlc* gene inactivated.

In the present invention, "L-amino acid producing bacterium" means a bacterium, which has an ability to accumulate L-amino acid in a medium, when the bacterium of the present invention is cultured in the medium. The L-amino acid producing ability may be imparted or enhanced by breeding. The term "L-amino acid producing bacterium" used herein also means a bacterium, which is able to produce and accumulate L-amino acid in a culture medium in amount larger than a wild type or parental strain of *E. coli*, such as *E. coli* K-12 strain.

The term “a bacterium belonging to the genus *Escherichia*” means that the bacterium is classified as the genus *Escherichia* according to the classification known to a person skilled in the microbiology. As examples of the microorganism belonging to the genus *Escherichia* used in the present invention, *Escherichia coli* (*E. coli*) can be mentioned.

The bacterium belonging to the genus *Escherichia* that can be used in the present invention is not particularly limited, however for example, bacteria described by Neidhardt, F.C. et al. (*Escherichia coli* and *Salmonella typhimurium*, American Society for Microbiology, Washington D.C., 1208, Table 1) can be mentioned.

The term “*mlc* gene is inactivated” means that the target gene is modified in the way that the modified gene encodes a mutant protein with decreased capacity for regulation or such gene encodes completely inactive protein. It is also possible, that the modified DNA region is unable to provide the natural expression of Mlc protein due to deletion of a part of the gene or modification of adjacent region of the gene.

Inactivation of *mlc* gene coding for the repressor of PTS brings an increasing the supply of carbon source, such as glucose, into the cell of L-amino acid producing bacterium.

The *mlc* gene codes for Mlc protein (406 amino acid residues), which is a global repressor of carbohydrate metabolism. The *mlc* gene (gi:16129552; numbers 1665368 to 1666588 in the GenBank accession number NC_000913.1) is located between *b1593* and *ynfL* genes on the chromosome of *E. coli* strain K-12.

Inactivation of the gene can be performed by conventional methods, such as mutagenesis treatment using UV irradiation or nitrosoguanidine (N-methyl-N'-nitro-N-nitrosoguanidine) treatment, site-directed mutagenesis, gene disruption using homologous recombination or/and insertion-deletion mutagenesis (Datsenko K.A. and Wanner B.L., Proc. Natl. Acad. Sci. USA, 2000, 97:12: 6640-45) also called as a “Red-driven integration”.

L- threonine producing bacterium.

As a bacterium of the present invention, which has to be modified to have *mlc* gene inactivated, the L-threonine producing bacteria may be used.

The bacterium of the present invention may be improved by enhancing the expression of one or more genes involved in the L-threonine biosynthesis. Such genes are exemplified by genes of L-threonine operon, i.e. *thr* operon, which preferably comprises the mutant *thrA* gene, which codes for aspartokinase homoserine dehydrogenase I resistant to feed back inhibition by threonine; the *thrB* gene, which codes for homoserine kinase; the *thrC* gene which codes for threonine synthase. Another preferred embodiment of the bacterium is modified to enhance the *rhtA* gene, which codes for putative transmembrane protein. Another preferred embodiment of the bacterium is modified to enhance the *aspC* gene, which codes for aspartate aminotransferase (aspartate transaminase) (Russian patent application No. 2002104983). The most preferred embodiment of the bacterium is modified to increase expression amount of *aspC* gene, the mutant *thrA* gene, the *thrB* gene, the *thrC* gene, the *rhtA* gene and modified to inactivate *mlc* gene.

As a parent strain, the L-threonine producing bacteria belonging to the genus *Escherichia* such as *E. coli* strain VKPM B-3996 (US Patent 5, 175, 107, US patent 5,705,371), *E. coli* strain NRRL-21593 (US Patent 5,939,307), *E. coli* strain FERM BP-3756 (US patent 5,474,918), *E. coli* strains FERM BP-3519 and FERM BP-3520 (US patent 5,376,538), *E. coli* strain MG442 (Gusyatiner et al., Genetika (in Russian), 14, 947-956 (1978)), *E. coli* strains VL643 and VL2055 (EP 1149911 A) and the like may be used.

The bacterium of the present invention can be obtained by inactivation of in the bacterium inherently having ability to produce L- amino acid. Alternatively, the bacterium of present invention can be obtained by imparting ability to produce L- amino acid to the bacterium already having *mlc* gene inactivated.

Methods for preparation of plasmid DNA, digestion and ligation of DNA, transformation, selection of an oligonucleotide as a primer and the like may be ordinary methods well known to

one skilled in the art. These methods are described, for instance, in Sambrook, J., Fritsch, E.F., and Maniatis, T., "Molecular Cloning A Laboratory Manual, Second Edition", Cold Spring Harbor Laboratory Press (1989).

2. Method of the present invention

The method of the present invention is a method for producing L-amino acid, which method comprises the steps of cultivating the bacterium of the present invention in a culture medium to produce and accumulate L-amino acid in the medium, and collecting L-amino acid from the medium. Concretely, the method of the present invention is a method for producing L-threonine, which method comprises the steps of cultivating the bacterium of the present invention in a culture medium to produce and accumulate L-threonine in the medium, and collecting L-threonine from the medium.

In the present invention, the cultivation, the collection and purification of L-amino acid from the medium and the like may be performed in a manner similar to the conventional fermentation method wherein an amino acid is produced using a bacterium.

A medium used for culture may be either a synthetic medium or a natural medium, so long as the medium includes a carbon source and a nitrogen source and minerals and, if necessary, appropriate amounts of nutrients which the bacterium requires for growth. The carbon source may include various carbohydrates such as glucose and sucrose, and various organic acids. Depending on the mode of assimilation of the used microorganism, alcohol including ethanol and glycerol may be used. As the nitrogen source, various ammonium salts such as ammonia and ammonium sulfate, other nitrogen compounds such as amines, a natural nitrogen source such as peptone, soybean-hydrolysate, and digested fermentative microorganism are used. As minerals, potassium monophosphate, magnesium sulfate, sodium chloride, ferrous sulfate, manganese sulfate, calcium chloride, and the like are used. As vitamins, thiamine, yeast extract and the like are used.

The cultivation is performed preferably under aerobic conditions such as a shaking culture, and stirring culture with aeration, at a temperature of 20 to 40 °C, preferably 30 to 38 °C. The pH of the culture is usually between 5 and 9, preferably between 6.5 and 7.2. The pH of the culture can be adjusted with ammonia, calcium carbonate, various acids, various bases, and buffers. Usually, a 1 to 5-day cultivation leads to the accumulation of the target L-amino acid in the liquid medium.

After cultivation, solids such as cells can be removed from the liquid medium by centrifugation or membrane filtration, and then L-amino acid can be collected and purified by ion-exchange, concentration and crystallization methods.

Brief Description of Drawings

Figure 1 shows relative position of the primers *mlcIL* and *mlcIR* on plasmid pACYC184 used for amplification of *cat* gene.

Figure 2 shows the construction of chromosomal DNA fragment comprising inactivated *mlc* gene.

Best Mode for Carrying out the Invention

The present invention will be more concretely explained below with reference to Examples.

Example 1. Construction the strain with inactivated *mlc* gene.

1. Deletion of *mlc* gene.

Deletion of *mlc* gene was constructed by means of the method firstly developed by Datsenko and Wanner (Proc. Natl. Acad. Sci. USA, 2000, 97(12), 6640-6645) called as a “Red-driven integration”. According to this procedure, the PCR primers *mlcIL* (SEQ ID NO: 1) and *mlcIR* (SEQ ID NO: 2) homologous to the both region adjacent to the *mlc* gene and gene conferring antibiotic resistance in the template plasmid were constructed. The plasmid pACYC184 (NBL Gene Sciences Ltd., UK) (GenBank/EMBL accession number X06403) was

used as a template in PCR reaction. Conditions for PCR were following: denaturation step for 3 min at 95 °C; profile for two first cycles: 1 min at 95 °C, 30 sec at 34 °C, 40 sec at 72 °C; profile for the last 30 cycles: 30 sec at 95 °C, 30 sec at 50 °C, 40 sec at 72 °C; final step: 5 min at 72 °C.

Obtained 935 bp PCR product (Fig. 1, SEQ ID NO: 3) was purified in agarose gel and used for electroporation of the *E. coli* strain MG1655, containing the plasmid pKD46 with temperature sensitive replication. The plasmid pKD46 (Datsenko and Wanner, Proc. Natl. Acad. Sci. USA, 2000, 97:12:6640-45) includes 2,154 nucleotides (31088-33241) DNA fragment of phage λ (GenBank accession No. J02459), containing genes of λ Red homologous recombination system (γ , β , exo genes) under control of arabinose-inducible P_{araB} promoter. The plasmid pKD46 is necessary for integration of the PCR product into chromosome of the strain MG1655.

Electrocompetent cells were prepared as follows: night culture of *E. coli* strain MG1655 grown at 30 °C in LB medium, supplemented with ampicillin (100 mg/l), was diluted in 100 times with 5 ml of SOB medium (Sambrook et al, “Molecular Cloning A Laboratory Manual, Second Edition”, Cold Spring Harbor Laboratory Press (1989)) with ampicillin and L-arabinose (1 mM). Obtained culture was grown with aeration at 30 °C to an OD_{600} of ≈ 0.6 and then made electrocompetent by concentrating 100-fold and washing three times with ice-cold deionized H_2O . Electroporation was performed using 70 μl of cells and ≈ 100 ng of PCR product. Cells after electroporation were incubated with 1 ml of SOC medium (Sambrook et al, “Molecular Cloning A Laboratory Manual, Second Edition”, Cold Spring Harbor Laboratory Press (1989)) at 37 °C for 2.5 h and after that plated onto L-agar and were growing at 37 °C to select Cm^{R} recombinants. Then to eliminate the pKD46 plasmid, 2 passages on L-agar with Cm at 42 °C were performed and the obtained colonies were tested for sensitivity to ampicillin.

2. Verification of *mlc* gene deletion by PCR.

The mutants, containing the deletion of *mlc* gene, marked with Cm resistance gene, were verified by PCR. Locus-specific primers *mlc*PL (SEQ ID NO: 4) and *mlc*PR (SEQ ID NO: 5)

were used in PCR for the verification. Conditions for PCR verification were following: denaturation step for 3 min at 94 °C; profile for the 30 cycles: 30 sec at 94 °C, 30 sec at 52 °C, 2 min at 72 °C; final step: 7 min at 72 °C. PCR product, obtained in the reaction with the cells of parental Mlc⁺ strain MG1655 as a template, was 1492 nucleotides in length (Fig.2, SEQ ID NO: 6). PCR product, obtained in the reaction with the cells of mutant MG1655 Δ mlc::cat strain as a template, was 1191 nucleotides in length (Fig.2, SEQ ID NO: 7).

3. Construction of threonine producing strain with inactivated *mlc* gene.

Threonine producing strain *E. coli* TDH7/pRT614 (VKPM B-5318, US patent 6,132,999) was transduced to Cm resistance by the standard procedure of P1 transduction (Sambrook et al, “Molecular Cloning A Laboratory Manual, Second Edition”, Cold Spring Harbor Laboratory Press (1989)). The strain MG1655 Δ mlc::cat was used as a donor. The resulted strain TDH7 Δ mlc::cat/pRT614 was verified by PCR to have Δ mlc::cat deletion by means of primers mlcPL (SEQ ID NO: 4) and mlcPR (SEQ ID NO: 5).

Example 2. Production of L-threonine by *E. coli* strain with inactivated *mlc* gene.

Both *E. coli* strain TDH7/pRT614 and TDH7 Δ mlc::cat/pRT614 was grown during 18-24 hours at 37 °C on L-agar plates containing streptomycin (50 µg/ml). Then one loop of the cells was transferred to 50 ml of L-broth of the following composition: trypton – 10g/l, yeast extract – 5 g/l, NaCl – 5 g/l. The cells (50 ml, OD₅₄₀ – 0.12 o.u.) grown at 37 °C within 4 hours on shaker (140 rpm) was used for seeding 450 ml of the medium for fermentation. The batch fermentation was performed in laboratory fermenter having a capacity of 1.0 l under aeration (1/1 vvm) with stirring at a speed of 1200 rpm at 39 °C. The pH value was maintained automatically at 6.6 using 8% ammonia liquor. The results are presented in Table 1.

The composition of the fermentation medium (g/l):

Glucose	100.0
NH ₄ Cl	1.75

KH ₂ PO ₄	1.0
MgSO ₄ x 7H ₂ O	0.8
FeSO ₄ x 7H ₂ O	0.01
MnSO ₄ x 5H ₂ O	0.01
Mameno(TN)	0.15
Betaine	1.0

Glucose and magnesium sulfate are sterilized separately. pH is adjusted to 6.6.

Table 1.

Strain	Amount of threonine, g/l	Yield, %	DCW, g/l	Cultivation time, h
TDH7/pRT614	12.6	13.3	16.6	22.8
TDH7Δ <i>mlc</i> ::cat/pRT614	16.9	18.2	16.8	21.7

As it is seen from Table 1, inactivation of the *mlc* gene improved the L-threonine accumulation by the L-threonine producing strain TDH7/pRT614.

SEQUENCE LISTING

<110> Ajinomoto-Genetika Research Institute

<120> METHOD FOR PRODUCING L-AMINO ACID USING BACTERIUM,
BELONGING TO THE GENUS ESCHERICHIA, LACKING ACTIVE MLC
GENE.

<140>

<141>

<160> 7

<170> PatentIn Ver. 2.1

<210> 1

<211> 64

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 1

agacgaatca acaaagaacc gttatacatc gcgtctatac ctgtgacgga agatcacttc gcag 64

<210> 2

<211> 63

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 2

cggagcgcga aaatataggg agtatgcggt ggttgcaatt acgccccgcc ctgccactca tcg 63

<210> 3

<211> 935

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cat gene
flanked by regions for integration

<400> 3

agacgaatca	acaaagaacc	gttatacatc	gcgtctatac	ctgtgacgga	agatcacttc	60
gcagaataaa	taaatcctgg	tgtccctggt	gataccggga	agccctgggc	caacttttgg	120
cgaaaatgag	acgttgatcg	gcacgtaaga	ggttccaact	ttcaccataa	tgaaataaga	180
tcactaccgg	gcgtatTTTT	tgagttatcg	agattttcag	gagctaagga	agctaaaatg	240
gagaaaaaaa	tactggata	taccaccggt	gatatatccc	aatggcatcg	taaagaacat	300
tttgaggcat	ttcagtcagt	tgctcaatgt	acctataacc	agaccgttca	gctggatatt	360
acggcctttt	taaagaccgt	aaagaaaaat	aagcacaagt	tttatccggc	ctttattcac	420
attcttgccc	gcctgatgaa	tgctcatccg	gaattccgta	tggaatgaa	agacgggtgag	480
ctggtgatat	gggatagtg	tcacccttgt	tacaccggtt	tccatgagca	aactgaaacg	540
ttttcatcgc	tctggagtga	ataccacgac	gattttccggc	agtttctaca	catatatctcg	600
caagatgtgg	cgtgttacgg	tgaaaaacctg	gcctatttcc	ctaaagggtt	tattgagaat	660
atgtttttcg	tctcagccaa	tccctggggt	agtttcacca	gttttgattt	aaacgtggcc	720
aatatggaca	acttcttcgc	cccgtttttc	accatgggca	aatattatac	gcaaggcgac	780
aaggtgctga	tgccgctggc	gattcaggtt	catcatgccg	tctgtgatgg	cttccatgtc	840
ggcagaatgc	ttaatgaatt	acaacagtac	tgcgatgagt	ggcagggcgg	ggcgtaattg	900

caaccaccgc atactcccta tattttcgcg ctccg

935

<210> 4

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 4

cagaagtgtc tgtaccggt

19

<210> 5

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: primer

<400> 5

aatgtgctgt taatcacatg c

21

<210> 6

<211> 1492

<212> DNA

<213> Escherichia coli

<400> 6

cagaagtgtc tgtaccggtg ataaagaaac gcttcagcat cactaactcc accgttatgc 60
ttcacaaata taaaccagga aaataattaa ccttgaaagt ctaagttatg ctttcctggc 120
ccaaattgag atagcgcaaa ttttggtaga acagttaaaa aatgttaacc ctgcaacaga 180
cgaatcaaca aagaaccgtt atacatcgcg tcttttacca gtgcagcgcc tgccatcggtg 240
ccctgggttag aaaactgagt actctcaacg ctgatgtgct gactatacgc aggaagggcc 300
tgctgacgga tgctgtctga gatgaccggg aagaggatat ctgccgcttt acttaacggt 360
gagccaatca gtattttttg tgggttaaat aaattcacca tgatggcaag aatgcgccg 420
acatgcgcgc ccaccccggt aatgatgtct tttgccagta gatcgccgcg caatgcgcc 480
tgacacaatg agtccacggt taacggttgt ccatgtaaca tcgagctcat ggattgatta 540
agacgcagct gtgccagctc aagaatactg tccacgctgg cgatggtttc gaggcagccg 600
tgattccgcg aataacagcg tttcccatc gggtcgacct gtgtgtggcc tatttccacg 660
agactactgc tgcctgcgtg tagcagatga ccatcggtaa tgacgccgcg cccacggtg 720
tgatcgataa ccacctgaat cacatcgcg gccccgcgtg aggcaccaa caaggcctct 780
gccatcgctc atgcgctgat atcatgctga atataaacg gaacgccggt atgctgctcc 840
agcgccctgc cgagcggcat ctcttttaca tctcgtaga acggcatgcg atgtacaata 900
ccattttccg tatcaataat tcccggcaag gttatggcaa tcgaagttag acgctcaagt 960
tttttctggt ggcggataaa aaactgatcg atatgggaaa taatacgatc cagcaatggc 1020
aagtcattct ttaacgccag ttcctgcgac tcttccacca ccagtttgct gctcagatcg 1080
cgcagagcaa ggaaaatctc cccgcgacta atgcgcagag aaagatagtg ccagggttca 1140
gtttcaacca ccagccccac cgccggacgg ccaagggttc ccgcttcttt gatttccagc 1200
tcttgacca ggtgtgcttc gagcatctca cggacaattt tagtgatact ggcaggagcc 1260
agttgcgcca gacgggaaag atcgatacgc gagactggac caagctgatc aatcaggcga 1320
taaaccgcgc ccgcgttggt ctgctttatt tgatcaatgt gccagcgtg gttttcagca 1380
accaccgcat actccctata ttttcgcgt ccgaaataat ctgtaggcta tgggtgaagca 1440
cttcaatacg tgcgtcaaaa tttttactta ggcattgtgat taacagcaca tt 1492

<210> 7

<211> 1196

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: delta mlc::cat
fragment

<400> 7

```
cagaagtgtc tgtaccggta ataaagaaac gcttcagcat cactaactcc accgttatgc 60
ttcacaaata taaaccagga aaataattaa ccttgaaagt ctaagttatg ctttcctggc 120
ccaaattgag atagcgcaaa ttttggtaga acagttaaaa aatgttaacc ctgcaacaga 180
cgaatcaaca aagaaccgtt atacatcgcg tctatacctg tgacggaaga tcacttcgca 240
gaataaataa atcctggtgt ccctgttgat accgggaagc cctgggcca cttttggcga 300
aaatgagacg ttgatcggca cgtaagaggt tccaactttc accataatga aataagatca 360
ctaccgggcg tattttttga gttatcgaga ttttcaggag ctaaggaagc taaaatggag 420
aaaaaaatca ctggatatac caccgttgat atatcccaat ggcatcgtaa agaacatttt 480
gaggcatttc agtcagttgc tcaatgtacc tataaccaga ccgttcagct ggatattacg 540
gcctttttta agaccgtaaa gaaaaataag cacaagtttt atccggcctt tattcacatt 600
cttgcccgcc tgatgaatgc tcatccggaa ttccgtatgg caatgaaaga cggtgagctg 660
gtgatatggg atagtgttca cccttggtac accgttttcc atgagcaaac tgaaacgttt 720
tcatcgctct ggagtgaata ccacgacgat ttccggcagt ttctacacat atattcgcaa 780
gatgtggcgt gttacggtga aaacctggcc tatttcctta aagggtttat tgagaatatg 840
tttttcgtct cagccaatcc ctgggtgagt ttcaccagtt ttgatttaaa cgtggccaat 900
atggacaact tcttcgcccc cgttttcacc atgggcaaat attatacgca aggcgacaag 960
gtgctgatgc cgctggcgat tcaggttcac catgccgtct gtgatggctt ccatgtcggc 1020
agaatgctta atgaattaca acagtactgc gatgagtggc agggcggggc gtaattgcaa 1080
ccaccgcata ctccctatat tttcgcgctc cgaaataatc tgtaggctat ggtgaagcac 1140
ttcaatacgt gtcgtcaaat ttttacttag gcatgtgatt aacagcacat t 1191
```


Fig. 1. Relative position of the primers *mlcIL* and *mlcIR* on plasmid *pACYC184*





Fig. 2. Construction of chromosomal DNA fragment comprising inactivated *mlc* gene



What is claimed is:

1. An L-amino acid producing bacterium belonging to the genus *Escherichia*, wherein the bacterium has been modified to have *mlc* gene inactivated.
2. The L-amino acid producing bacterium, wherein L-amino acid is L-threonine.
3. A method for producing L-amino acid, which method comprises the steps of:
 - cultivating the bacterium according to claim 1 in a medium to produce and accumulate L-amino acid in the medium, and
 - collecting L-amino acid from the medium.
4. The method according to claim 3, wherein L-amino acid is L-threonine.
5. The method according to claim 4, wherein the bacterium has been modified to have enhanced expression of L-threonine operon.

Abstract of disclosure

There is provided a method for producing L-amino acid, such as L-threonine, using bacterium belonging the genus *Escherichia* with inactivated *mlc* gene coding for global regulator of carbohydrate metabolism.